

## Purification and TEM of Afp and Its Variants

Daria Rybakova<sup>1\*</sup>, Alok K. Mitra<sup>2</sup> and Mark R. H. Hurs<sup>1</sup>

<sup>1</sup>Innovative Farm Systems, AgResearch, Lincoln Research Centre, Christchurch, New Zealand;

<sup>2</sup>School of Biological Sciences, University of Auckland, Auckland, New Zealand

\*For correspondence: [dariaryb@gmail.com](mailto:dariaryb@gmail.com)

**[Abstract]** The *Serratia entomophila* antifeeding prophage (Afp), forms a phage-tail-like particle (tailocin) that causes cessation of feeding activity of the New Zealand grass grub, *Costelytra zealandica*. Here, we describe the more detailed purification protocol for Afp particles and its variants which is based on the procedure described in our original publication (Rybakova *et al.*, 2013). The purification procedure includes inducing *Escherichia coli* (*E. coli*) cells harbouring *afp* genes under arabinose-inducible promoter for 24 h. The cells are harvested and sonicated on ice followed by DNase treatment and centrifugation. The supernatant is then filter sterilised and applied to the size exclusion chromatography (SEC) column. The fractions containing Afp or its variants are pooled and ultracentrifuged. The supernatant is removed and the transparent pellet is resuspended in the residual buffer. The procedure results in Afp particles or variants thereof that are approximately 70% pure. The Afp particles are negatively stained and visualized using Transmission electron microscopy (TEM).

### **Materials and Reagents**

*Note: All solutions, unless otherwise specified, were prepared with distilled deionised Millipore-microfiltered (MilliQ) water, autoclaved for 15 min at 1.1 kg/cm<sup>2</sup> at 121 °C and stored at room temperature. If antibiotics or supplements were to be added, the medium was cooled to a temperature of 50-60 °C before addition of antibiotic or supplement solution. The following antibiotics were used at stated concentrations (ug/ml): ampicillin, 100; chloramphenicol, 30; and spectinomycin, 100.*

1. *E. coli* strains containing *afp* genes under arabinose promoter

*Note: Please refer to the original manuscript (Rybakova *et al.*, 2013) for strains and antibiotic resistance description.*

2. LB-agar (Merck KGaA, catalog number: VM459383236)
3. DNase (Roche Diagnostics, catalog number: 04716728001)
4. 0.7 % filtered (pore size 0.2 um) uranyl acetate diluted in sterile water (pH 5.0) (Electron Microscopy Sciences, catalog number: 22400)

5. Sephacryl S-400 HR resin (GE,  
[http://www.gelifesciences.com/webapp/wcs/stores/servlet/catalog/en/GELifeSciences/products/AlternativeProductStructure\\_17395/17060901](http://www.gelifesciences.com/webapp/wcs/stores/servlet/catalog/en/GELifeSciences/products/AlternativeProductStructure_17395/17060901))
6. 20% L-Arabinose (Acros organics) (Thermo Fisher Scientific, catalog number: AC104985000) (see Recipes)
7. 0.4x Luria-Bertani broth (LB) (Life Technologies, catalog number: 12795-027) (see Recipes)
8. 0.2% sodium azide (Sigma-Aldrich, catalog number: S8032) (see Recipes)
9. 250 mM TBS buffer (see Recipes)
10. TM buffer (see Recipes)

## **Equipment**

1. Spectrophotometer
2. Orbital mixer incubator (Raytek Corporation)
3. 1.7 ml microcentrifuge tubes
4. Eppendorf centrifuges
5. Sanyo soniprep 150 sonicator (18 Ω) (Panasonic Corporation)
6. 0.45-μm-pore-size cellulose acetate filters (Advantec MFS, catalog number: 03CP045AN)
7. 50 ml BD Falcon Tubes
8. Sephacryl S-400 HR beads (GE Healthcare, catalog number: 17-0609-10) for size exclusion chromatography (SEC)
9. The SEC column with a bed volume of 1.5 by 46 cm (Bio-Rad Laboratories, model: 737-4155)
10. 1.5 x 50 cm (D x L) column (Farmacia)
11. BioLogic LP system (Bio-Rad Laboratories, model: 731-8350)
12. Beckman coulter Optima TM L-100K ultracentrifuge (Beckman Coulter, model: 392050)
13. Polaron E5100 SEM coating unit  
*Note: This unit is from before 2000 and is no longer available, however a similar unit that is equivalent can be found at <http://www.quorumtech.com/products/glow-discharge-system-free-standing.html>*
14. Whatman #1 filter paper (Whatman)
15. Plastic-coated 200-mesh grid (ProSciTech, catalog number: GCU200)
16. Morgagni 268D TEM (FEI)
11. Megapixel III digital camera  
*Note: The catalog numbers and model numbers are not important for reproducibility.*

12. Polypropylen tubes used for the size exclusion chromatography (polypropylene 12 x 75 mm, 5 ml) (Labserv, catalog number: LBS504N)
  13. Polypropylen tubes for ultracentrifugation (polycarbonate centrifuge bottles with cap assembly 16 x 76 mm, capacity 10.4 ml) (Beckman Coulter, catalog number: 355603)
  14. Falcon tubes (Thermo Fisher Scientific, catalog number 14-432-22)
  1. Protect microorganism preservation System (Technical Service Consultants) for preservation of *E. coli* cells at -80 °C
- Note: When required, a single inoculated bead was removed from the vial and dropped into a LB broth containing the appropriate antibiotic.*

## **Procedure**

*Note: Centrifugation steps were performed at 4 °C unless otherwise noted.*

### A. Afp purification protocol

1. A single inoculating bead (Protect Microorganism Preservation System) from the -80 °C stock of an appropriate Afp or its derivatives bearing *E. coli* strain is used to inoculate 3 ml of LB with the respective antibiotics.
2. The culture is incubated overnight at 37 °C with agitation at 250 rpm (overnight culture).
3. 1 ml of the overnight culture is used to inoculate 50 ml of fresh LB with appropriate antibiotic/s.
4. The 50 ml of the cell culture is grown at 37 °C with agitation to an OD<sub>600</sub> of 1.0.  
*Note: OD<sub>600</sub> measurements are taken from the cell cultures diluted with the LB media at ratios of 1:10. A cuvette with LB medium is used as a blank.*
5. The cells are transferred into sterile falcon tubes (50 ml) and harvested by centrifugation at 2,268 x g for 10 min at room temperature.
6. The cell pellets are resuspended in fresh 50 ml 0.4x LB broth with respective antibiotic/s.  
*Note: Initially 1 ml of the 0.4x LB broth is added to each pellet and the pellet is resuspended by vigorous tapping on the tube. Then the tube is filled up to 50 ml with 0.4x LB broth and the content of the tube is transferred to the same 50 ml flask as the culture was grown. Antibiotics are added directly to the flask.*
7. 500 µl of steril filtered 20% arabinose (final concentration 0.2%) is added to each 50 ml culture to induce the protein expression.  
*Note: After arabinose is added, avoid shaking the cultures.*
8. The cultures are incubated for a further 24 h at 22 °C at 40 rpm.
9. The induced cells are harvested by centrifugation for 10 min at 2,268 x g.  
*Note: Higher centrifugation speed (4,629 x g for 10 min) may be necessary if pellet is too loose.*

10. The pellet from each 50 ml culture is re-suspended at 4 °C in 0.8 ml pre-chilled TM buffer.
11. The resultant approximately 0.9 ml re-suspensions are transferred into 1.7 ml microcentrifuge tubes and subjected to three 20-sec rounds of sonication using a Sanyo Soniprep 150 Sonicator (18 Ω). At least 1 min of incubation on ice is included between sonication steps.
12. One µl (10 units) of the DNase is added and the sample is left at room temperature for 10 min.
13. Following the DNase incubation step the suspension is centrifuged at 16,000 x g for 10 min at 4 °C.
14. The pellets are discarded and clear supernatants containing soluble recombinant protein are filtered through 0.45-µm-pore-size filters.

*Notes:*

- a. *If the samples need to be stored for 1-2 days, 0.02% (final concentration) NaN<sub>3</sub> is added to each tube.*
  - b. *Storage longer than 2 days was not tested; however for a better reproducibility it is recommended to proceed with the purification immediately after this step.*
15. The resin is packed into a 1.5 x 50 cm (D x L) column according to the manufacturers' instructions, resulting in a column bed 1.5 cm in diameter and 46 cm in height.  
*Note: The SEC column can be reused.*
  16. The column is attached to the BioLogic LP System and equilibrated with 150-200 ml TBS at a flow rate of 2 ml/min.
  17. Protein solution is then loaded and eluted in a 65 ml of buffer at 0.5 ml/min at room temperature. Elution of protein was monitored by absorbance at 280 nm with the typical OD<sub>280</sub> values of 0.01-0.2.

SEC protocol:

Period of time (min)	Flow rate (ml/ min)	Action
0-40	0.5	Discard
40-130	0.5	Collect 1.5 min per fraction (0.75 ml each fraction)
130-200	0.5	Divert to waste


18. The 1.5 ml fractions from the elution peak (approximately 75-90 min retention time) are collected and pooled as follows: fraction numbers 18-27 (pool I); 28-31 (pool II); 32-46 (pool III).

*Note: The pool II usually contains the Afp or its derivatives with the least amount of cellular contaminants, while pool I and III contain less concentrated Afp particles with higher amounts of cellular contaminants. However, due to slight variations in column filling density, the fractions to be pooled have to be determined experimentally each time the column is repoured.*

19. The pooled SEC fractions containing Afp or its derivatives are centrifuged at 151,139 x g for one hour and 10 min at 4 °C in a Beckman Optima TM L-100K ultracentrifuge (fixed angle).

*Notes:*


- a. *It is helpful to label the outside of the tube where the pellet is likely to precipitate.*
  - b. *The tube after the centrifugation has to be handled very carefully. Proceed to the next step immediately after the ultracentrifugation step to avoid disturbance of the pellet.*
20. After the centrifugation the supernatant is carefully removed using a pipette without disturbing the pellet but leaving approximately 150-200 µl that was left enabling the later resuspension of the pellet (Figure 1).



**Figure 1. Removing supernatant after ultracentrifugation.** The supernatant is removed using a pipette; approximately 150-300 µl covering the invisible pellet is left behind.

21. The tube is incubated for 30-60 min at 22 °C at 40 rpm, tilted with the labelled side down (Figure 2) in order to gently solubilise the pellet.

Residual buffer (150-200 ul with the pellet)



**Figure 2. Gentle resuspension of the pellet.** The tube is placed as shown in the figure and gently agitated for 30-60 min.

22. Following incubation, the often transparent retentate containing particles of interest is resuspended in residual buffer (usually approximately 150-200  $\mu$ l) by pipetting up and down.
23. The resuspended pellet is transferred to the 1.7 ml Eppendorf tube and centrifuged for 30 sec at 16,000  $\times$  *g*. The pellet is discarded and supernatant is transferred to a new eppendorf tube with 0.02% sodium azide (final concentration) added. This preparation is used for visualizing Afp particles or its variants using transmission electron microscopy or for other applications.


*Note: Afp particles of approximately 70% purity can be purified using this method.*

#### B. Negative staining and TEM of Afp and its variants

1. The grids are charged by using glow discharging following manufacturer's instructions no longer than 15 min before applying a sample.
2. To visualise Afp and its variants, 3  $\mu$ l of the re-suspended pellet is applied to a freshly glow-discharged plastic-coated 200-mesh grid.
3. The sample is allowed to settle for 60 sec.
4. Excess fluid is wicked off using Whatman #1 filter paper and then stained for 45 sec with a 0.7% filtered uranyl acetate (pH 5.0).

*Note: 0.5-1.0% can also be used.*

5. The grid is left to air dry on a filter paper for 5-15 min.
6. Grids are examined in a Morgagni 268D TEM at a nominal magnification of 70,000-180,000 operated at 80 KeV following manufacturer's instructions. The images are captured using a Megapixel III digital camera. Several examples of the Afp images are shown in Figure 3.



**Figure 3. Examples of the uncropped Afp images taken at nominal magnifications 89,000 (A) and 110,000 (B)**

### Recipes

1. 20% L-Arabinose (5 ml stock solution)  
Mix 1 g Arabinose with 5 ml dH<sub>2</sub>O
2. 0.4x LB (50 ml)  
Mix 20 ml LB and 30 ml dH<sub>2</sub>O
3. 0.2% sodium azide (10 ml stock solution)  
Mix 0.02 g sodium azide with 10 ml dH<sub>2</sub>O
4. 250 mM TBS (pH 7.4) (500 ml 10x stock)  
Mix 40 g NaCl  
1 g KCl

- 2.42 g Tris base
  - 16.5 g Tris-HCl
  - With 490 ml dH<sub>2</sub>O
  - Ajust pH to 7.5 with HCl (approximately 1 ml 6 M HCl)
  - Add dH<sub>2</sub>O to 500 ml
  - Use in 1:10 dilution (25 mM)
  - Filter sterilize (0.45 µm) before using for SEC
5. TM buffer
- Mix 2.52 g Tris-HCl with 3.25 g MgCl<sub>2</sub>
  - Add dH<sub>2</sub>O to 800 ml and adjust pH to 7
  - Autoclave and sterile filter (0.45 µm)

### **Acknowledgments**

The protocol was adapted from Rybakova *et al.* (2013). We are grateful to the following colleagues from AgResearch: Dr Duane Harland for help with TEM, Sandra Jones for assisting in protein purification, Chikako van Koten for statistical assessments and Stefan Clerens for LC-MS/MS. Sam Yu (Izon Science, Christchurch, New Zealand) is thanked for the assessment of Afp particles. This research was funded by Grant C10X0804 of the New Economy Research Fund (NERF), administered by the New Zealand Foundation for Research, Science and Technology.

### **References**

1. Rybakova, D., Radjainia, M., Turner, A., Sen, A., Mitra, A. K. and Hurst, M. R. (2013). [Role of antifeeding prophage \(Afp\) protein Afp16 in terminating the length of the Afp tailocin and stabilizing its sheath.](#) *Mol Microbiol* 89(4): 702-714.