

## Determination of Survival of Wildtype and Mutant *Escherichia coli* in Soil

Yinka Somorin and Conor O'Byrne\*

Microbiology, School of Natural Sciences, National University of Ireland, Galway, Ireland

\*For correspondence: [conor.obyrne@nuigalway.ie](mailto:conor.obyrne@nuigalway.ie)

**[Abstract]** *E. coli* resides in the gastrointestinal tract of humans and other warm-blooded animals but recent studies have shown that *E. coli* can persist and grow in various external environments including soil. The general stress response regulator, RpoS, helps *E. coli* overcome various stresses, however its role in soil survival was unknown. This soil survival assay protocol was developed and used to determine the role of the general stress response regulator, RpoS, in the survival of *E. coli* in soil. Using this soil survival assay, we demonstrated that RpoS was important for the survival of *E. coli* in soil. This protocol describes the development of the soil survival assay especially the recovery of *E. coli* inoculated into soil and can be adapted to allow further investigations into the survival of other bacteria in soil.

**Keywords:** Soil survival, *Escherichia coli*, RpoS, Environmental persistence, Bacteria recovery

**[Background]** *Escherichia coli* is a Gram-negative, facultative anaerobe, belonging to the Enterobacteriaceae family, which inhabits the intestinal tract of humans, warm-blooded animals and reptiles (Berg, 1996; Gordon and Cowling, 2003). It can be transferred through water and sediments via faeces and is used as an indicator of faecal contamination in drinking and recreational water. The use of *E. coli* as a faecal indicator is based, at least in part, on the assumption that it exists transiently outside of the host gastrointestinal tract (Ishii and Sadowsky, 2008) and does not survive for a long time in the external environment. However, several studies have isolated *E. coli* from various natural environments such as municipal wastewater, freshwater, beach water, beach sand and soils (Jiménez *et al.*, 1989; Brennan *et al.*, 2010; Chiang *et al.*, 2011; Byappanahalli *et al.*, 2012; Zhi *et al.*, 2016). The capacity of these *E. coli* strains to survive for long periods of time and grow in the external environment raises questions about the validity of its continued use as indicator of water quality (Brennan *et al.*, 2010). To understand the genetic mechanism underlying the survival and persistence of *E. coli* in soil, we developed a soil survival assay to evaluate the role of the different genetic factors on soil survival. We investigated the role of the general stress response regulator, RpoS, in the survival of long-term soil persistent *E. coli* in soil. The ability of the *rpoS* mutant (COB583 $\Delta$ *rpoS*) to survive in soil was compared with the wildtype strain (COB583) and RpoS was demonstrated to be important for the survival and long-term persistence of *E. coli* in soil (Somorin *et al.*, 2016). Here, we present the detailed protocol for the soil survival assay and the recovery of *E. coli* from soil.

## **Materials and Reagents**

*Note: All reagents used were from the specified manufacturers (catalog numbers indicated). Nonetheless, the same reagents from different manufacturers are expected to produce similar results.*

1. Spatula
2. Weighing boat (Sparks Lab Supplies, catalog number: BAL1822)
3. Gloves (Sparks Lab Supplies, catalog number: SAF5534X20)
4. Sterile micropipette tips (100-1,000 µl) (Abdos Labtech, catalog number: P10102)
5. Sterile micropipette tips (2-200 µl) (Abdos Labtech, catalog number: P10101)
6. Microcentrifuge tube (1.5 ml) (Abdos Labtech, catalog number: P10202)
7. Centrifuge tube (15 ml graduated) (Abdos Labtech, catalog number: P10402)
8. Cuvette (LP ITALIANA, catalog number: 112117)
9. 2 mm laboratory test sieve (B5410/1986, Endecotts, catalog number: 200SIW2.00)
10. 96-well plate (TC microwell 96U) (Thermo Fisher Scientific, Thermo Scientific™, catalog number: 163320)
11. 90 mm Petri dishes (Abdos Labtech, catalog number: P10901)
12. Silty-Loam soil
13. *E. coli* strains
  - a. *E. coli* COB583 (Somorin *et al.*, 2016)
  - b. *E. coli* COB583Δ*rpoS* (Somorin *et al.*, 2016)

*Note: E. coli stock cultures were made in LB with 7% Dimethyl sulfoxide (v/v; Sigma-Aldrich) and kept at -80 °C pending use.*
14. Luria-Bertani (LB) broth (Sigma-Aldrich, catalog number: L3022-1KG)
15. Agar No. 2 (Lab M, catalog number: MC006)
16. MacConkey agar (Sigma-Aldrich, catalog number: M7408)
17. Distilled water (Milli-Q) (EMD Millipore)
18. Phosphate buffered saline (PBS) (Sigma-Aldrich, catalog number: P4417-100TAB)
19. LB broth (see Recipes)
20. LB agar plates (see Recipes)
21. PBS buffer (see Recipes)
22. MacConkey agar plates (see Recipes)

## **Equipment**


1. 250 ml conical flask (VWR, catalog number: 214-1132)
2. -80 °C freezer
3. Styrofoam 15 ml tube holder
4. Bunsen burner

5. Orbital shaker (Gallenkamp)
6. Discovery comfort multichannel pipette (20-200 µl; HTL)
7. Discovery comfort multichannel pipette (5-20 µl; HTL)
8. Nichipet EX pipette (200-1,000 µl; Nichiryo)
9. Nichipet EX Pipetman classic pipettes (20-200 µl; Nichiryo)
10. Vortex mixer (Reax top; Heidolph)
11. Biomate 3 spectrophotometer (Thermo Fisher Scientific, catalog number: 335904)  
*Note: This product has been discontinued.*
12. Weighing scale (Sartorius, catalog number: BL120S)  
*Note: This product has been discontinued.*
13. Centrifuge (Eppendorf, model: 5418)
14. Labo autoclave (SANYO, model: MLS-3020U)

## **Procedure**

- A. Measurement of cell concentration by optical density (OD<sub>600 nm</sub>)
  1. Streak out *Escherichia coli* COB583 wildtype and the *rpoS* deletion mutant (COB583Δ*rpoS*) from the stock culture kept in the -80 °C freezer onto LB agar (see Recipes) around a Bunsen burner and incubate them overnight (~16 h) at 37 °C.
  2. Inoculate one colony of each strain with a disposable sterile inoculating loop into 10 ml LB in a sterile conical flask around a Bunsen burner and incubate them at 37 °C for 6 h. Perform this step in triplicate for each strain.
  3. From the 6 h culture, take 100 µl and add it to 900 µl LB and determine the OD<sub>600 nm</sub>.
  4. Determine the volume of the 6 h culture required to give a starting OD<sub>600 nm</sub> of 0.05 in a 25 ml LB as follows:  
$$C_1V_1 = C_2V_2$$
$$C_1 = \text{OD}_{600 \text{ nm}} \text{ of the 6 h culture; } V_1 = \text{volume of the 6 h culture}$$
$$C_2 = \text{final OD}_{600 \text{ nm}} \text{ required; } V_2 = \text{final volume required}$$
$$V_1 = 0.05 \times 25 \text{ ml}/C_1$$
  5. Make the cultures with starting OD<sub>600 nm</sub> of 0.05 in 25 ml LB for each strain and their respective replicates in sterile 250 ml conical flasks and incubate the flasks overnight on a shaker at 37 °C.
  6. On the following day, dilute the overnight cultures 1:10 in LB and determine the OD<sub>600 nm</sub>.
  7. Pipette 1 ml of each overnight culture into two separate sterile 1.5 ml tubes for each strain and their respective replicates.
  8. Centrifuge the cultures at 9,000 x g for 10 min at room temperature (23-25 °C) and remove the supernatants with sterile pipette.
  9. Wash the cell pellets by re-suspending them in sterile PBS (see Recipes) and centrifuge at 9,000 x g for 10 min. Repeat the washing step two more times.


10. Resuspend the washed cell pellets in 1 ml of sterile PBS by pipetting up and down and vortexing and dilute 1:2; 1:5; 1:10; 1:20; 1:100; and 1:1,000 in sterile PBS to make 1 ml in sterile microcentrifuge tubes.
11. Transfer each diluted cell suspension into a cuvette and determine the OD<sub>600 nm</sub> of the diluted cell suspensions of the *E. coli* strains and their respective replicates in a spectrophotometer three times and determine the mean value.
12. Make serial dilutions of each cell suspension and spot 10 µl onto LB plates in triplicate and incubate the plates at 37 °C overnight.
13. Count the colonies and use the numbers derived to estimate the cell count (CFU ml<sup>-1</sup>) and determine the mean cell count from the triplicate experiments.
14. Plot the mean cell count against the mean OD<sub>600 nm</sub> of the respective dilutions of each strain in three independent experiments to obtain calibration curves from which desired cell numbers can be obtained. The standard curve of graph of the cell count versus OD<sub>600 nm</sub> is presented in Figure 1.



**Figure 1. Standard curve of cell numbers versus OD<sub>600 nm</sub> of COB583 (A) and COB583ΔrpoS (B)**

## B. Development of soil survival assay

1. Inoculate one colony each of COB583 and COB583 $\Delta$ rpoS into 25 ml LB in triplicate and incubate overnight at 37 °C.
2. Dilute the resulting cultures 1:10 in LB and determine the OD<sub>600 nm</sub>.
3. Add 1 ml of the overnight culture to two sterile Eppendorf tubes for each replicate of the two strains.
4. Wash the cultures as described above (steps A8 and A9).
5. Pool the resulting cell pellets and resuspend in 2 ml of sterile PBS in a 15 ml tube.
6. Make 1:10 dilution (1 ml) of the each washed cell suspension to determine the OD<sub>600 nm</sub>.
7. Based on the OD<sub>600 nm</sub>, prepare a normalised cell suspension with OD<sub>600 nm</sub> equivalent to 2 x 10<sup>8</sup> CFU ml<sup>-1</sup> from each replicate.
8. Make serial dilution of each normalised bacterial suspension and spot 10  $\mu$ l of each dilution on LB agar in triplicate. Allow the spots to air-dry and incubate the plates overnight at 37 °C. Count the colonies and determine the mean cell counts.
9. Sieve the soil to be used with 2 mm laboratory test sieve and weigh 1 g of the sieved soil into a 15 ml tube.
10. Take 50  $\mu$ l of each normalised bacterial suspension for each strain and inoculate it into the soil (to give ~1 x 10<sup>7</sup> CFU g<sup>-1</sup> of soil). The uninoculated control should have 50  $\mu$ l of sterile PBS added to it. The experiment was set up in triplicate for each strain.
11. Cover the tubes with the lid and invert them 10 times by hand for proper mixing.
12. Leave the tubes for about 10 min at room temperature (23-25 °C).
13. Perform the recovery of the inoculated *E. coli* strains by adding 2 ml of sterile PBS to each 15 ml tube. Cover the tube with its lid, invert it three times and immediately vortex for 20 sec twice.
14. Leave the tube on the workbench and allow the larger soil particles in the resulting soil slurry to settle out (2-10 min, depending on soil texture) (Figure 2).




**Figure 2. Experimental set-up for the soil survival assay with the soil slurry just after mixing and after settling out**

15. Take 20  $\mu\text{l}$  from the supernatant of each soil slurry, place them in separate wells in a 96-well plate and make serial dilution (up to  $10^{-6}$ ) from them.
16. Take 10  $\mu\text{l}$  of each dilution and spot onto MacConkey agar plates (see Recipes) in triplicate, and incubate plates overnight at 37 °C.
17. Count the colonies and use the numbers derived to estimate the mean recovered cell count (CFU  $\text{ml}^{-1}$ ) from the triplicate experiments. The percentage recovery of cells was determined.

### Data analysis

Cell count for each replicate is represented by the mean cell counts obtained from the three spots in each replicate. The mean and standard deviation of the cell counts from the normalised cell suspension (inoculum) and recovered cell counts were obtained from three independent replicates of each strain. The percentage of recovery ranges from 30-79% and the recovered cells from this procedure is presented in Figure 3.

$$\text{Percentage Recovery (\%)} = \frac{\text{Recovered cell count}}{\text{Inoculum cell count}} \times 100$$



**Figure 3. Recovery of *Escherichia coli* from soil.** Percentage recovery is provided above the recovered cells.

### Notes

1. The optical density of the inoculum should be confirmed to be accurate before inoculating into soil.
2. Care should be taken when inoculating the soil to ensure the pipette tips do not contact the soil directly, as it prevents the release of the entire content of the pipette tip.
3. Care should be taken when spotting the cell suspension on agar plates so as not to puncture the agar with the pipette tips. This could vary the cell numbers arising from each spot.

4. The soil to be used should be checked and confirmed not to contain *E. coli* prior to the soil survival assay so that recovered *E. coli* will only be attributable to inoculated *E. coli*. Serial dilutions of 1 g of the soil sieved through a 2 mm mesh should be plated on MacConkey agar and incubated at 37 °C to determine this.
5. For long-term soil survival experiments, the tubes should be slightly capped to allow air exchange and incubated statically at the desired temperature. Multiple tubes should be set up for experiments requiring sampling at different time points since inoculated soils are destructively sampled (*i.e.*, PBS was added and could not be reused at another time point) for *E. coli* recovery.
6. No significant difference was seen when soil slurry was left to settle for 2 and 10 min before aliquots were taken for serial dilution and plating for cell count determination.

### **Recipes**

1. LB broth  
Add 20 g LB powder to 1 L of distilled water and autoclave
2. LB agar plates  
Add 20 g LB powder and 15 g agar No. 2 powder to 1 L of distilled water and autoclave. When cool to touch (~ 45 °C), pour about 25 ml of the molten agar into Petri dishes and allow to solidify
3. PBS buffer  
Dissolve one phosphate-buffered saline tablet (Sigma-Aldrich) in 200 ml of distilled water and autoclave, giving a PBS buffer (pH 7.4)
4. MacConkey agar plates  
Add 50 g MacConkey agar powder to 1 L of distilled water and autoclave. When cool to touch (~45 °C), pour about 25 ml of the molten agar into Petri dishes and allow to solidify

### **Acknowledgments**

The protocol was adapted from a previously published work (Somorin *et al.*, 2016). The authors thank Camila Thorne for her help and advice with setting up the soil survival assay. This work was supported by an NUI Galway College of Science Ph.D. Fellowship and Thomas Crawford Hayes research award to Y.S.

### **References**

1. Anderson, K. L., Whitlock, J. E. and Harwood, V. J. (2005). [Persistence and differential survival of fecal indicator bacteria in subtropical waters and sediments](#). *Appl Environ Microbiol* 71(6): 3041-3048.
2. Berg, R. D. (1996). [The indigenous gastrointestinal microflora](#). *Trends Microbiol* 4(11): 430-435.

3. Brennan, F. P., O'Flaherty, V., Kramers, G., Grant, J. and Richards, K. G. (2010). [Long-term persistence and leaching of \*Escherichia coli\* in temperate maritime soils](#). *Appl Environ Microbiol* 76(5): 1449-1455.
4. Byappanahalli, M. N., Whitman, R. L., Shively, D. A., Sadowsky, M. J. and Ishii, S. (2006). [Population structure, persistence, and seasonality of autochthonous \*Escherichia coli\* in temperate, coastal forest soil from a Great Lakes watershed](#). *Environ Microbiol* 8(3): 504-513.
5. Byappanahalli, M. N., Yan, T., Hamilton, M. J., Ishii, S., Fujioka, R. S., Whitman, R. L. and Sadowsky, M. J. (2012). [The population structure of \*Escherichia coli\* isolated from subtropical and temperate soils](#). *Sci Total Environ* 417-418: 273-279.
6. Chiang, S. M., Dong, T., Edge, T. A. and Schellhorn, H. E. (2011). [Phenotypic diversity caused by differential RpoS activity among environmental \*Escherichia coli\* isolates](#). *Appl Environ Microbiol* 77(22): 7915-7923.
7. Gordon, D. M. and Cowling, A. (2003). [The distribution and genetic structure of \*Escherichia coli\* in Australian vertebrates: host and geographic effects](#). *Microbiology* 149(Pt 12): 3575-3586.
8. Jimenez, L., Muniz, I., Toranzos, G. A. and Hazen, T. C. (1989). [Survival and activity of \*Salmonella typhimurium\* and \*Escherichia coli\* in tropical freshwater](#). *J Appl Bacteriol* 67(1): 61-69.
9. Ishii, S., Ksoll, W. B., Hicks, R. E. and Sadowsky, M. J. (2006). [Presence and growth of naturalized \*Escherichia coli\* in temperate soils from Lake Superior watersheds](#). *Appl Environ Microbiol* 72(1): 612-621.
10. Ishii, S. and Sadowsky, M. J. (2008). [Escherichia coli in the environment: Implications for water quality and human health](#). *Microbes Environ* 23(2): 101-108.
11. Somorin, Y., Abram, F., Brennan, F. and O'Byrne, C. (2016). [The general stress response is conserved in long-term soil-persistent strains of \*Escherichia coli\*](#). *Appl Environ Microbiol* 82(15): 4628-4640.
12. Zhi, S., Banting, G., Li, Q., Edge, T. A., Topp, E., Sokurenko, M., Scott, C., Braithwaite, S., Ruecker, N. J., Yasui, Y., McAllister, T., Chui, L. and Neumann, N. F. (2016). [Evidence of naturalized stress-tolerant strains of \*Escherichia coli\* in municipal wastewater treatment plants](#). *Appl Environ Microbiol* 82(18): 5505-5518.