

## Immunofluorescence (Especially for Cells Growing on a Coverglass)

Hui Zhu\*

Department of Genetics, Stanford University, Stanford, CA, USA

\*For correspondence: [huizhu@stanford.edu](mailto:huizhu@stanford.edu)

**[Abstract]** If an antibody for your protein of interest is available, immunofluorescence is a useful method to detect the localization and relative abundance of the protein by using a fluorescence microscope. Immunofluorescence can be used in combination with other, non-antibody methods of fluorescence staining, for example, the use of DAPI to label DNA. This protocol describes setting up an immunofluorescence experiment using cells grown on a coverglass.

### Materials and Reagents

1. Primary antibodies: self-produce or commercially order  
*Note: For self-produce primary antibodies, optimize antibody concentration at 1-50  $\mu\text{g/ml}$ ; Try 5  $\mu\text{g/ml}$  as a starting point. For commercial primary antibodies, usually 1:1,000 dilution as a starting point, and then adjust the working concentration according to the preliminary data.*
2. Secondary antibodies (below are the most common secondary antibodies):
  - a. Goat anti-rabbit IgG (Life Technologies, Molecular Probes®/Alexa Fluor® 488, catalog number: A-11008)
  - b. Goat anti-rabbit IgG (Life Technologies, Molecular Probes®/Alexa Fluor® 594, catalog number: A-11012)
  - c. Goat anti-mouse IgG (Life Technologies, Molecular Probes®/Alexa Fluor® 488, catalog number: A-11001)
  - d. Goat anti-mouse IgG (Life Technologies, Molecular Probes®/Alexa Fluor® 594, catalog number: A-11005)
  - e. Goat anti-human IgG (Life Technologies, Molecular Probes®/Alexa Fluor® 488, catalog number: A-11013)
  - f. Goat anti-human IgG (Life Technologies, Molecular Probes®/Alexa Fluor® 594, catalog number: A-11014)*Note: We usually dilute commercial secondary antibodies at 1:200 as a starting point. Make sure to spin down the stock solution of secondary antibody before dilution to prevent precipitated fluorophores from causing high background. We usually aliquot secondary antibody and stored at -20 °C.*

3. 4', 6-diamidino-2-phenylindole dihydrochloride (DAPI) (Boeringer Mannheim, 236 276) (make 1 mg/ml stock, 1,000x)
4. Paraformaldehyde (16% solution) (Electron Microscopy Sciences, catalog number: 15170)
5. Methanol (Thermo Fisher Scientific, catalog number: BP 1105-4)
6. Coverglasses (12 mm) (Thermo Fisher Scientific, catalog number: 12-545-82)
7. Poly-L-lysine (MW 150,000-300,000) (Sigma-Aldrich, catalog number: P1399)
8. Fluormount-G™ (mounting solution) (SouthernBiotech, catalog number: 0100-01)
9. Nitric acid
10. Hydrochloric acid
11. Liquid nitrogen
12. PBS-BT solution (see Recipes)
13. Normal goat serum (The Jackson Laboratory, catalog number: 005-000-001) (see Recipes)

### **Equipment**

1. Glass beaker
2. Petri dishes
3. UV-irradiate
4. 24-well plate
5. 6-well plate
6. 60 mm dish
7. Parafilm

### **Procedure**

#### A. Preparing coverglasses

1. Make up 100 ml acid solution in a large glass beaker in the hood.  
*Note: The acid solution is made of 2 parts of nitric acid and 1 part of HCl, and the color is orange-red.*
2. Put the 12 mm coverglasses into the acid solution one by one so that the coverglasses are evenly washed in the acid.
3. Let the coverglasses sit in the acid for 2 h or overnight swirling occasionally.
4. Decant acid solution into another large glass beaker; neutralize acid with NaOH pellets.  
*Note: With NaOH, the solution will 'boil' vigorously, so be careful.*
5. Wash coverglasses with ddH<sub>2</sub>O until pH goes up to 7.0.

6. Add sufficient 500 µg/ml poly-L-lysine solution to cover all coverglasses, and sit for 2 h with occasionally swirling.
7. Decant poly-L-lysine solution (solution can be used repeatedly).
8. Wash coverglasses 3 times with ddH<sub>2</sub>O.
9. Dry each coverglass individually in Petri dishes, avoiding coverglasses stuck to the surface.

*Note: If coverglasses are stuck to the surface of a Petri dish, they can be easily removed by adding a small amount of liquid nitrogen.*

10. After dry, put all the coverglasses into a Petri dish and UV-irradiate overnight before first use.

*Note: For coverglasses that are stored for a long time, re-irradiate for the sake of sterility.*

#### B. Growing cells on coverglasses

11. Split cells onto tissue culture dishes containing coverglasses or chambered slides.

*Notes:*

- a. *We usually put one coverglass in one well of the 24-well plate; or no more than 2 coverglasses in one well of the 6-well plate, or no more than 5 coverglasses in a 60 mm dish.*
  - b. *Pipette up and down or shake the dish to make sure cells are not concentrating in the center of the dish well. Make sure there are no air bubbles between the coverglasses and the tissue culture dish.*
12. Grow cells to 70-100% confluency.

#### C. Fixation and permeabilization of the cells

13. Transfer the coverglasses or chambered slides into another tissue plate containing sufficient methanol (-20 °C stock), fix for 10 min or a couple of weeks.

Alternately, transfer cells to another plate including PBS. Discard PBS, adding 200 µl PBS plus 20 µl 16% paraformaldehyde. Shaking the plate for 15 min at room temperature (RT).

*Notes:*

- a. *Transfer to methanol directly, no need to do the PBS wash.*
- b. *16% paraformaldehyde needs to be stored at -20 °C, the final working concentration is 1.6% here.*
- c. *Formaldehyde preserves antigens by crosslinking the proteins, it has the advantage of preserving most subcellular antigens in their proper localization and GFP. Methanol precipitates antigens by dehydrating cells, it is useful for observing*

*cytoskeletal elements such as microtubules, actin, and other associated structures, but it destroys GFP.*

14. Carefully transfer the coverglasses or chambered slides from the plates and place cell side up onto secured Parafilm. Wash the coverglasses or chambered slides with 100  $\mu$ l PBS immediately after transfer, never dry the cells.
15. Add 100  $\mu$ l PBS-BT solution to the coverglasses or chambered slides, let sit for 30 min at RT to permeablize and block cells.

*Note: For stringent block, 4-6% normal goat serum (NGS) can be used.*

#### D. Staining and mounting cells

16. Incubate cells in 40  $\mu$ l primary antibody (1  $\mu$ g/ml final primary antibody concentration, dilute in PBS-BT) for 30 min at RT.
17. Rinse with PBS-BT twice, and then wash with 100  $\mu$ l PBS-BT twice, 5 min each.
18. Cells were incubated in 40  $\mu$ l secondary antibody (1  $\mu$ g/ml final secondary antibody concentration, dilute in PBS-BT) for 30 min at RT.
19. Rinse with PBS-BT twice, wash with PBS-BT once, 5 min, and then wash with PBS, 5 min.
20. Incubate cells in 40  $\mu$ l DAPI (1  $\mu$ g/ml final concentration, 1:1,000 dilute in PBS) for 2 min, and then wash with PBS once.
21. Add 5-10  $\mu$ l mounting solution to a clean microscope slide for each coverglass, place stained coverglass cell side down onto mounting solution from one edge; allow mounting solution to cover the entire surface of the coverglass, avoiding air bubbles.
22. Let the mounting solution dry and self-seal for 30 min at RT.

#### Recipes

1. PBS-BT solution
  - 10 ml 10x PBS
  - 3 g BSA (to 3%)
  - 1 ml 10% Triton X-100 (to 0.1%)
  - 1 ml 5% NaN<sub>3</sub>
  - ddH<sub>2</sub>O to 100 ml
  - Stored at 4 °C
2. Normal goat serum
  - Make 4-6% solution in PBS

## **Acknowledgments**

This protocol was modified from an immunoprecipitation protocol developed in the laboratory of Dr. Guowei Fang (Department of Biology, Stanford University, Stanford, CA, USA). The protocol was originally developed Dr. Jim Wong. This work was supported by a Burroughs-Wellcome Career Award in Biomedical Research (G.F.) and by grants from National Institutes of Health (GM062852 to G.F.).

## **References**

1. Wong, J. and Fang, G. (2006). [HURP controls spindle dynamics to promote proper interkinetochore tension and efficient kinetochore capture.](#) *J Cell Biol* 173(6): 879-891.
2. Zhu, H., Coppinger, J. A., Jang, C. Y., Yates, J. R., 3rd and Fang, G. (2008). [FAM29A promotes microtubule amplification via recruitment of the NEDD1-gamma-tubulin complex to the mitotic spindle.](#) *J Cell Biol* 183(5): 835-848.
3. Zhu, H., Fang, K. and Fang, G. (2009). [FAM29A, a target of Plk1 regulation, controls the partitioning of NEDD1 between the mitotic spindle and the centrosomes.](#) *J Cell Sci* 122(Pt 15): 2750-2759.